

## Amended Abstract

**Objectives:** To assess the potency of CEM-101, a fluoroketolide in clinical development, against strains observed to be resistant (R) to other agents in the same class. Reports have documented ketolide-R (telithromycin [TEL]) species worldwide, most recently *S. pyogenes* from Europe. CEM-101 was tested against a collection of 43 TEL-R  $\beta$ -haemolytic streptococci (BHS).

**Methods:** A total of 53 (1.3%) BHS were identified among 3,958 in the SENTRY Antimicrobial Surveillance Program (2003-2006) that were TEL-R (MIC,  $\geq 2$  mg/L). 43 strains (36 Group A, 1 Group C, 6 Group G) were available for testing, from 20 hospitals in Europe (31 strains), North America (11) and Latin America (1). Susceptibility (S) testing used CLSI broth microdilution methods and 3 strains were erythromycin (ERY)-R, clindamycin (CC)-S requiring D-test. Nine comparison agents were tested (4 in Table).

**Results:** The potency of CEM-101 against each BHS serogroup was the same with an overall MIC<sub>50</sub> and MIC<sub>90</sub> of 0.12 and 0.5 mg/L, respectively. CEM-101 activity was 32-fold (MIC<sub>50</sub> comparisons) greater than TEL. All strains were ERY-R, but 100% S to quinupristin/dalfopristin (Q/D). Three CC-S strains (*S. pyogenes*) were D-test (+) and 2 had (+) induction of R to CEM-101. The S rates for other comparators were: penicillin, tetracycline, ceftriaxone, amoxicillin/clavulanate, and levofloxacin (100.0%); and tetracycline (46.8%).

Table. MIC distributions for CEM-101 and MLS<sub>B</sub>-ketolide comparison agents.

Antimicrobial	Occurrences at MIC (mg/L):									
	$\leq 0.015$	0.03	0.06	0.12	0.25	0.5	1	2	4	>4
CEM-101	4	0	1	18	10	6	4	0	0	0
Telithromycin	0	0	0	0	0	0	0	8	16	19
Erythromycin	0	0	0	0	0	0	0	-	43	
Clindamycin	-	-	-	2	1	0	0	-	40	
Q/D <sup>a</sup>	-	-	-	37	6	0	0	-	0	

a. Q/D = quinupristin/dalfopristin

**Conclusions:** CEM-101 remained active against all TEL-R (MIC,  $\geq 2$  mg/L) BHS with all MICs at  $\leq 1$  mg/L (MIC<sub>50</sub>, 0.12 mg/L). Highest occurrence of TEL-R strains was in Europe (greatest in Italy). CEM-101 warrants further development for infections caused by BHS.

## Introduction

CEM-101 is a novel fluoroketolide selected as a candidate for parenteral and oral therapy of community-acquired respiratory tract (CA-RTI) and for other infections where macrolides are indicated. Initial in vitro studies indicated a potency comparable to or superior to telithromycin (a ketolide), erythromycin, azithromycin and clarithromycin, as well as activity against Gram-positive strains having documented resistances to macrolides or lincosamides. CEM-101 activity is generally focused against Gram-positive pathogens, but also possesses measurable potencies versus fastidious Gram-negative species (*Haemophilus*, *Moraxella*), some Enterobacteriaceae (*Salmonella*, *Shigella*) and pathogens causing various sexual transmitted diseases (STD; *Neisseria gonorrhoeae*, *Chlamydia trachomatis*).

In this presentation, we report CEM-101 activity measured by reference Clinical and Laboratory Standards Institute (CLSI) methods when testing organisms recently (2008) recognized as resistant to telithromycin.

These  $\beta$ -haemolytic streptococci have emerged mainly in Europe and possess telithromycin MIC values of  $\geq 2$  mg/L for 5.2% of isolates in 2004-2005. *S. pyogenes* (Group A) has been the dominant species sampled in Western Europe with this phenotype.

The collection of the SENTRY Antimicrobial Surveillance Program was searched for these ketolide-non-susceptible strains and tested against CEM-101 to assess the potential advantage for the newer agent in the same structural antimicrobial class.

## Materials and Methods

### Organisms collected:

All organisms tested were obtained from patients in the United States (USA), Europe or Latin America. A total of 53 (1.3%)  $\beta$ -haemolytic streptococci (3,958 isolates) had elevated telithromycin MIC values at  $\geq 2$  mg/L, the level consistent with "non-susceptibility" by USA-FDA and CLSI breakpoint criteria when applied to streptococci such as *S. pneumoniae* (CLSI M07-A8, 2009 and M100-S19, 2009).

Forty-three of these strains were available for further study, from 20 hospitals in Europe (31 strains), USA (11 strains), and Latin America (one strain). The most common species were *S. pyogenes* (Group A; 36 strains), Group C (one strain) and Group G (six strains).

### Susceptibility testing:

Broth microdilution (96-well) panels were produced by TREK Diagnostics (Cleveland, Ohio, USA) to be used with 2.5-5% lysed horse blood supplement to test fastidious streptococci. More than 30 antimicrobials were tested (10 shown here) including investigational agents such as CEM-101. All tests used the CLSI M07-A8 (2009) method and quality control (QC)/breakpoints guidelines from CLSI M100-S19 (2009). Tested QC strains included *S. aureus* ATCC 29213, *E. faecalis* ATCC 29212, *S. pneumoniae* ATCC 49619 and *H. influenzae* ATCC 49247 and 49766. All QC MIC results were within published ranges.

Those organisms having clindamycin susceptibility, but resistance to erythromycin were tested for inducible lincosamide resistance by the CLSI D-test per M100-S19 (2009) method. This was only used for three strains (all *S. pyogenes*).

## Results

- Telithromycin-non-susceptible  $\beta$ -haemolytic streptococci were more commonly *S. pyogenes* (2.4%) when compared to all strains tested (1.3%) during 2002-2006 (3,958 isolates); see Table 1.

Table 1. Telithromycin MIC distributions for *S. pyogenes* and all  $\beta$ -haemolytic streptococci (BHS) tested in the SENTRY Antimicrobial Surveillance Program (2002-2006).

Organism (no. tested)	Cum. % inhibited at MIC (mg/L)				
	$\leq 0.5$	1	2	4	>4
<i>S. pyogenes</i> (1,631)	96.6	<u>97.6</u> <sup>a</sup>	98.1	99.1	100.0
All BHS (3,958)	97.9	<u>98.7</u> <sup>a</sup>	98.9	99.3	100.0

a. Underlined % indicates susceptible population (CLSI breakpoint for *S. pneumoniae*).

- Table 2 shows that CEM-101 had MIC values at  $\leq 1$  mg/L for all of the 43 available telithromycin-non-susceptible  $\beta$ -haemolytic streptococcal strains (MIC<sub>50</sub>, 0.12 mg/L; Table 3).

Table 2. Activity of CEM-101 and nine comparative agents tested against 43  $\beta$ -haemolytic streptococci with elevated telithromycin MIC values ( $\geq 2$  mg/L).

Antimicrobial agent	No. of occurrences at MIC (mg/L)								
	$\leq 0.03$	0.06	0.12	0.25	0.5	1	2	4	>4
CEM-101	4	1	18	10	6	4	0	0	0
Telithromycin	0	0	0	0	0	<u>0</u> <sup>a</sup>	8	16	19
Erythromycin	0	0	0	<u>0</u>	0	0	0	- <sup>b</sup>	43
Clindamycin	-	-	-	<u>2</u>	1	0	0	-	40
Q/D <sup>c</sup>	-	-	-	37	6	<u>0</u>	0	-	0
Penicillin	42	1	<u>0</u>	0	0	0	0	0	0
Ceftriaxone	-	-	-	43	<u>0</u>	0	0	0	0
Levofloxacin	-	-	-	-	39	3	<u>1</u>	0	0
Tetracycline	-	-	-	-	-	-	<u>21</u>	0	22
TMP/SMX <sup>c</sup>	-	-	-	-	43	0	0	0	0

a. Underline value at breakpoint concentration (CLSI). TMP/SMX has no breakpoint criteria.

b. - = untested concentration.

c. Q/D = quinupristin/dalfopristin and TMP/SMX = trimethoprim/sulfamethoxazole.

- Antimicrobials remaining active against these telithromycin refractory strains were (MIC<sub>90</sub>/% susceptible): Q/D (0.5/100.0), penicillin ( $\leq 0.03/100.0$ ), ceftriaxone ( $\leq 0.25/100.0$ ), and levofloxacin (1/100.0), see Table 2. Only 46.8% of these strains were tetracycline-susceptible.
- Macrolides and clindamycin were not active against the test strains and the three clindamycin-susceptible strains were D-test-positive (data not shown), i.e., inducible clindamycin-resistant.
- Table 3 illustrates that the total population of telithromycin-non-susceptible strains and the *S. pyogenes* subset had identical MIC<sub>50/90</sub> results. Group G isolates (six total) had a slightly higher MIC<sub>50</sub> (0.5 mg/L; data not shown).

Table 3. Comparative activity of CEM-101 tested against *S. pyogenes* and all  $\beta$ -haemolytic streptococci (BHS) with telithromycin resistance.

Organism (no. tested)	MIC (mg/L)			
	50%	90%	Range	% $\leq 1$ mg/L
<i>S. pyogenes</i> (36)	0.12	0.5	$\leq 0.015$ -1	100.0
All BHS (43)	0.12	0.5	$\leq 0.015$ -1	100.0

## Conclusions

- Telithromycin resistance has emerged in  $\beta$ -haemolytic streptococci, most commonly in Europe (72.1% of occurrences by region) and among *S. pyogenes* (83.7% of occurrences by species).
- A new fluoroketolide, CEM-101, was 32-fold more potent than telithromycin and all MIC values were  $\leq 1$  mg/L, that breakpoint used for telithromycin when testing other streptococci such as *S. pneumoniae*.
- CEM-101 activity against  $\beta$ -haemolytic streptococci appears to be superior to telithromycin and could provide a wider and sustained coverage of tested species. Further development is warranted, especially in endemic areas (Europe) of ketolide resistance (*erm* [B] and 23S rRNA mutations).

## Selected References

- Bemer-Melchior P, Juvin ME, Tassin S, Bryskier A, Schito GC and Drugeon HB (2000). In vitro activity of the new ketolide telithromycin compared with those of macrolides against *Streptococcus pyogenes*: influences of resistance mechanisms and methodological factors. *Antimicrob. Agents Chemother.* 44:2999-3002.
- Clinical and Laboratory Standards Institute (2009). *M07-A8. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; approved standard - eighth edition.* Wayne, PA., CLSI.
- Clinical and Laboratory Standards Institute (2009). *M100-S19. Performance standards for antimicrobial susceptibility testing. 19th informational supplement.* Wayne, PA. CLSI.
- Douthwaite S, Jalava J and Jakobsen L (2005). Ketolide resistance in *Streptococcus pyogenes* correlates with the degree of rRNA dimethylation by Erm. *Mol. Microbiol.* 58:613-622.
- Farrell DJ, Morrissey I, Bakker S and Felmingham D (2002). Molecular characterization of macrolide resistance mechanisms among *Streptococcus pneumoniae* and *Streptococcus pyogenes* isolated from the PROTEKT 1999-2000 study. *J. Antimicrob. Chemother.* 50 Suppl S1:39-47.
- Farrell DJ, Shackcloth J, Barbadora KA and Green MD (2006). *Streptococcus pyogenes* isolates with high-level macrolide resistance and reduced susceptibility to telithromycin associated with 23S rRNA mutations. *Antimicrob. Agents Chemother.* 50:817-818.
- Grivea IN, Al-Lahham A, Katopodis GD, Syrogiannopoulos GA and Reinert RR (2006). Resistance to erythromycin and telithromycin in *Streptococcus pyogenes* isolates obtained between 1999 and 2002 from Greek children with tonsillopharyngitis: phenotypic and genotypic analysis. *Antimicrob. Agents Chemother.* 50:256-261.
- Malhotra-Kumar S, Lammens C, Chapelle S, Wijdooghe M, Piessens J, Van Herck K, et al. (2005). Macrolide- and telithromycin-resistant *Streptococcus pyogenes*, Belgium, 1999-2003. *Emerg. Infect. Dis.* 11:939-942.
- Nagai K, Appelbaum PC, Davies TA, Kelly LM, Hoellman DB, Andrasciv AT, et al. (2002). Susceptibility to telithromycin in 1,011 *Streptococcus pyogenes* isolates from 10 central and Eastern European countries. *Antimicrob. Agents Chemother.* 46:546-549.
- Reinert RR, Luttkien R, Sutcliffe JA, Tait-Kamradt A, Cil MY, Schorn HM, et al. (2004). Clonal relatedness of erythromycin-resistant *Streptococcus pyogenes* isolates in Germany. *Antimicrob. Agents Chemother.* 48:1369-1373.
- Richter SS, Heilmann KP, Dohrn CL, Beekmann SE, Riah F, Garcia-de-Lomas J, et al. (2008). Increasing telithromycin resistance among *Streptococcus pyogenes* in Europe. *J. Antimicrob. Chemother.* 61:603-611.
- Tamayo J, Perez-Trallero E, Gomez-Garcas JL and Alos JI (2005). Resistance to macrolides, clindamycin and telithromycin in *Streptococcus pyogenes* isolated in Spain during 2004. *J. Antimicrob. Chemother.* 56:780-782.